



Ruptured Dilated Right Coronary Sinus of Valsalva with Ventricular Septal Defect

Dr Ma. Nina Castillo

Training and Medical Center, La Union 2500, Philippines

Abstract:

Ventricular septal defects account for 10 percent of congenital heart diseases in adults. They can either occur alone, or in combination with other cardiac anomalies. Ventricular septal defects, and more commonly aortic valve regurgitation and bicuspid aortic valve, are frequently associated with ruptured sinus of Valsalva aneurysms. Rupture of the aneurysm is the most common complication, which commonly protrude and rupture into the right ventricle or right atrium.

We present the case of a 27-year old male who was admitted because of lightheadedness and dizziness. The patient had a history of a heart murmur since childhood and was diagnosed with a congenital heart disease. 2D-echocardiography with Doppler studies revealed a dilated right



Biography:

Health Social Scientist (Sociology, Clinical Epidemiology, Global Health); and Human Resources Development & Management Consultant

Publications:

1. Evaluating the Mechanical Properties of Admixed Blended Cement Pastes and Estimating its Kinetics of Hydration by Different Techniques
2. Genetic Diversity Using Random Amplified Polymorphic DNA (RAPD) Analysis for *Aspergillus niger* isolates
3. Au-Ag-Cu nanoparticles alloys showed antifungal activity against the antibiotics-resistant *Candida albicans*
4. Induce mutations for Bavistin resistance in *Trichoderma harzianum* by UV-irradiation
5. Biliary Sludge. Analysis of a Clinical Case

[2nd World cardiology Experts Meeting September 21-22, 2020](#)

Abstract Citation: [Dr Ma. Nina Castillo, Ruptured Dilated Right Coronary Sinus of Valsalva with Ventricular Septal Defect, WORLD CARDIOLOGY SUMMIT 2020, 2nd World cardiology Experts Meeting, September 21-22, 2020](#)